

Hot Pepper



MOMBEY F1

Jalapeno



HIGHLIGHTS:

Compact Vegetation

■ Uniform & Attractive Fruit Color

High Set

Pickling & Fresh Consumption

Slot	Maturity	Cropping	Fruit Size	Fruit Color	Pungency
Spring	Average 80 Days	Open Field	Average 8x3 cm	Yellow	Strong

^{*} Days to Maturity are from transplanting date. They may vary due to weather fluctuation, growing season and location.

Disease Package

HR:

IR:

 ${\it Disease\ Package\ Resistance\ Keys:\ HR=High\ Resistance;\ IR=Intermediate\ Resistance.}$

 $\textbf{\textit{TMV-}} To bacco\ \textit{Mosaic Virus}; \textbf{\textit{Tm-}} To bamo\ \textit{Viruses}; \textbf{\textit{TSWV-}} To mato\ \textit{Spotted\ Wilt\ Virus}; \textbf{\textit{PepMoV-}} Pepper\ \textit{Mottle\ Virus}; \textbf{\textit{PVY-}} Potato\ \textit{Virus\ Y}; \textbf{\textit{Rs-}} Bacterial\ \textit{Wilt\ Xcv-} Bacterial\ \textit{Spot\ Lt-} Leveillula\ taurica; \textbf{\textit{Pe-Phtophthora}} blight; \textbf{\textit{N-}} Root\ \textit{Knot.}$

Disclaime

All published data on this website are indicative only and based on crop performance observed in our trial stations. Performance may vary from one geographical area to another and from year to year as local growing habits, soil, and weathe conditions may vary. Distributors are urged to run local trials to determine the abiotic and biotic (pathoge strains, etc.) effects on the growin hybrid(s).

Read on the above statement, we give no varranty for crop performance perfaining to the informance and the province of t