



Greenhouse Cucumber



CASPER F1

**Beit Alpha
Parthenocarpic**



HIGHLIGHTS:

Balanced Vegetation

Main Stem Bearing Habit

Short Internodes

Consistent Fruit Quality

Slot	Maturity	Flowering Mode	L/D Ratio	Skin Color	Skin Texture
Spring Autumn	Average 35 Days	Semi Multi	Average 19 x 3 cm	Glossy	Fine Ribbing

** Days to Maturity are from transplanting date. They may vary due to weather fluctuation, growing season and location.*

Disease Package

HR: Px

IR: CMV

*Disease Package Resistance Keys: **HR** = High Resistance, **IR** = Intermediate Resistance*

*Disease Code: **CMV** - Cucumber mosaic virus, **CVYV** - Cucumber vein yellowing virus, **CYSDV** - Cucurbit yellow stunting disorder virus, **WMV** - Watermelon mosaic virus, **ZYMV** - Zucchini yellow mosaic virus, **PRSV** - Papaya ringspot virus, **Pcu** - Downy mildew, **Px** - Powdery mildew, **Ccu** - Scab*

Disclaimer:
All published data on this website are indicative only and based on crop performance observed in our trial stations. Performance may vary from one geographical area to another and from year to year as local growing habits, soil, and weather conditions may vary. Distributors are urged to run local trials to determine the abiotic and biotic (pathogen strains, etc.) effects on the grown hybrid(s).
Based on the above statement, we give no warranty for crop performance pertaining to the information given; nor do we accept any liability for any loss, direct or ensuing, that may result from any cause.